

Noise from wind turbines and risk of cardiovascular disease

The number and size of wind turbines has increased markedly during the last decades. Although research has consistently found that traffic noise increases the risk for cardiovascular disease (1-4), potential effects of wind turbine noise on risk for cardiovascular disease are virtually unexplored.

The aim of this project is to investigate if exposure to wind turbine noise is associated with an increased risk for cardiovascular disease using unique Danish registers. We will investigate if:


- acute exposure to wind turbine noise triggers myocardial infarction and stroke
- long-term exposure to wind turbine noise increases the risk of cardiovascular disease

BACKGROUND

Hearing is a permanent process essential for human survival and communication. However, the human organism is not able to shut off noise and therefore noise can have a number of unwanted effects, such as reduction of fidelity of communication, interference with cognitive processes and disturbance of sleep.

Figure 1 shows two pathways through which noise exposure can affect human health: directly (a) and indirectly via sleep disturbance (b + c) (5).

Noise acts as a stressor and, according to the general stress model, provokes a typical stress response, including hyperactivity of the sympathetic autonomic nervous system and activation of the hypothalamus–pituitary– adrenal axis, resulting in vasoconstriction, increased blood pressure, increased heart rate and high levels of the glucocorticoid cortisol (6, 7). Also, acute stress has been found to affect the immune system with increased levels of circulating inflammatory factors such as IL-6 and IL-1 β (8).



Exposure to noise during night-time is thought particularly hazardous (5). Studies have shown that noise at night causes a full range of sleep disturbance from minor unconscious autonomic perturbations, such as sleep stage changes and body movements, to full awakening (9-11). Sleep disturbance has been associated with changes in physiological parameters, such as metabolic and endocrine function (12, 13) and altered immune defense, such as increased levels of immunoglobulins (14, 15).

Longer term exposure to noise from road traffic and airports has consistently been found to increase the risk of cardiovascular disease: hypertension, myocardial infarction (MI) and stroke (1-4). However, noise immission from wind turbines and traffic are different, which highlights the need for research of potential cardiovascular health effects of wind turbine noise. Firstly, wind turbine noise levels are generally low compared to traffic noise levels in urban settings. In Denmark wind turbines emits max. 44 dB whereas for road traffic noise approximately one third of all Danish dwelling are exposed to more than 58 dB (L_{den}). Also, as the sound power level from a wind turbine depends on the wind velocity,

the immission levels vary irregularly (16), whereas traffic noise usually are more constant and predictable. In a recent study, Janssen et al. found that wind turbine noise induced a higher proportion of annoyed residents than traffic noise does at comparable sound levels (16). Also, annoyance due to wind turbine noise was found at lower noise levels than traffic noise. However, the study also showed that annoyance was lower among residents who received economical benefit from wind turbines (16).

Wind turbine noise has been suggested to be associated with sleep disturbances, stress and general health, although results are inconsistent (17-22). All these studies are cross-sectional and uses self-reported data on sleep, stress and health. Also, they are generally based on a small number of exposed persons. Some of these studies indicate that associations between wind turbine noise and sleep, stress and health are present only among annoyed people whereas there seems to be no relationship between modeled/measured wind turbine noise and sleep disturbances, stress and health among people that report not be annoyed by noise from a nearby wind turbine(s) (18, 21). One study has investigated associations between modeled wind turbine noise and self-reported cardiovascular disease and they found no associations (19). However, the study was cross-sectional, based on self-reported cardiovascular disease, the number of persons with as well as the type of cardiovascular disease was not reported and results only adjusted for age and sex, and therefore this hypothesis should be evaluated in a larger, more solid design, ideally a prospective study which is considered the best design for studying health effects of long-term exposure.

Acute effects of noise on cardiovascular events are virtually unexplored, although noise-induced risk factors including increased blood pressure and inflammation are believed to be important in the trigger of a stroke and a MI (23, 24). A large day-to-day variation in noise exposure is needed to investigate acute effects. In contrast to traffic noise, wind turbines noise is associated with considerable day-to-day variation, providing a unique opportunity to investigate whether noise exposure trigger a cardiovascular event immediately after exposure.

In Denmark we have unique opportunities to investigate whether wind turbine noise affects the risk of cardiovascular disease because Denmark has been a pioneer in setting up wind turbines and therefore we can follow the health effects of wind turbines over many years (Figure 2), and secondly because Denmark has unique national registers that enables us to trace the addresses of all Danish citizens since 1971 and all hospital admissions for cardiovascular disease since 1977.




Figure 2 Growth in number of wind turbines in Denmark (modified from energinet.dk)

METHODS

Exposure assessment

The Danish Energy Agency and Energinet.dk have established a register of all Danish wind turbines. This register is updated monthly, and contains information on all wind turbines back to 1980 including the exact location of each wind turbine (geographical coordinate), date of grid connection, cancellation date for decommissioned turbines, manufacture, type, hub height and diameter of rotor blades. The registry contains data for approximately 5,000 wind turbines in operation and 2,500 cancelled wind turbines.

There is a clear association between wind velocity and the noise emitted from a wind turbine which is characteristic and available for each type of wind turbine (25). We will use these data to develop a model for noise immission from all types of wind turbines at all wind velocities. Based on these modeled noise levels for each turbine we will identify all dwellings within the immission area of wind turbine noise using geographical information system (GIS), and the noise level at each dwelling will then be calculated using the distance between the wind turbine and the dwelling. It will in many cases be necessary to calculate the noise contributions from several wind turbines, which will then be summed for each dwelling.

Nord2000 is a highly accurate method for calculation of noise, verified by controlled measurements including measurements of wind turbine noise (26). As calculations with Nord2000 are very time consuming and costly, we will use a simplified version of the method described in details in "Vindmøllebekendtgørelsen" from the Danish EPA (27). We consider this method to be well fitted for the considerable number of calculations needed for this project. In addition, we will correct all modeled values for the attenuation that occurs when sound propagates in headwind (see appendix 1).

Simulated wind speed and direction time series at each wind turbine location in Denmark back to 1980 will be produced based on the mesoscale wind analysis method developed by DTU Wind Energy and used in many studies (28-30). These data will be provided at a spatial resolution of approximately 5 km and downscaled to each required location at a 1-hour or 3-hour resolution and at different levels, including 10 m above ground and hub height. The wind analysis method makes use of the advanced Weather Research and Forecasting (WRF) model (31) and a dynamic downscaling atmospheric reanalysis technique (32).

Sub-study 1: Day-to-day variation in wind turbine noise and risk for cardiovascular disease

Previous studies on traffic noise and risk of cardiovascular disease have focused on long-term noise exposure. One reason is that there is little day-to-day variation in exposure to traffic noise. In contrast, exposure to noise from wind turbines is associated with considerable day-to-day variation, which gives us a unique opportunity to investigate whether noise exposure can lead to MI or stroke immediately after exposure.

Study population and design

Based on modeled noise immission from all types of wind turbines (see exposure section) we will identify all residential addresses within the immission area using GIS and the official Danish address database (Den Offentlige Informationsserver (OIS), www.ois.dk). Subsequently, we will identify all persons who have lived at those addresses in the period 1980-2012 by linking the addresses with the Danish civil registration system, thereby retrieving their unique personal identification number (33). We will include all persons above 25 years of age at start of exposure (set up of new turbine or moving into a noise immission area) who were exposed to noise from wind turbines in more than one year in the period from 1980 to 2012. We estimate that this will sum up to approximately 10,000–15,000 persons. This estimate is based on the following: 1) 7,500 wind turbines in Denmark (total of turbines in operation and cancelled), 2) for most single wind turbines there will be one dwelling just below the noise limit value, as this determines the dimensions of the turbine and the required distance to dwellings, 3) for smaller groups of wind turbines (typically 3-5) we estimate that 2-3 dwellings will be exposed just below the noise limit value, 4) for large wind farms with up to 100 turbines we estimate that 3-10 dwellings will be exposed just below the noise limit value, and 5) for both single and multiple wind turbine areas we expect that in many cases there will also be dwellings within the immission area but at lower exposure than the limit value.

We will identify the persons among this exposed population that have been hospitalized at least once with stroke and/or MI during the maximum of 32 years we follow them. We estimate that this will sum up to approximately 2,000 relevant

hospitalizations during the study period based on information on number of yearly events of the two diseases (www.hjerteforeningen.dk). We use stroke and MI as endpoints as we have previously found road traffic noise to increase the risk for these diseases (3, 4). Information on hospitalization for stroke (International Classification of Disease (ICD) 10: I61, I63 and I64) and MI (ICD10: I21.0-I21.9) in the period 1980-2012 for all exposed persons will then be collected from the Danish National Patient Registry (34), and information on death will be collected from the Danish register of causes of death (35) using the personal identification number. For all exposed cases with MI/stroke we will calculate equivalent continuous A-weighted wind turbine sound pressure level (L_{Aeq}) at their residence for the each day (L_d ; 07:00–19:00 h), evening (L_e ; 19:00–22:00 h) and night (L_n ; 22:00–07:00 h) as described in the exposure section. The low time resolution enables us to investigate health effects of wind turbine noise at different times of the day, which is of great interest as one hypothesis is that exposure during night is particular hazardous (5).

We will use a case-crossover design, where we estimate the relationship between day-to-day variability in wind turbine noise at each address and day-to-day variation in hospitalizations for cardiovascular disease (stroke and MI) among people exposed to wind turbines noise. In a case-crossover study the noise exposure of a case person on the day of hospitalization (and a few days before) is compared with noise exposure during a control period. The advantage of the design is that each person is their own control and differences in socio-economic status and lifestyle factors are accounted for via the study design.

Daily variations in air pollution and outdoor temperature could potentially confound the results. The majority of all Danish wind turbines are placed in rural settings where air pollution levels are mainly determined by the regional background contribution. Department of Environmental Science at Aarhus University has historical daily measurements on regional air pollution background levels as well as temperature which will be included in this study.

Statistical analyses

We will estimate risk using conditional logistic regression and compare the noise exposure on the day a person is hospitalized with stroke/MI (lag 0) with the noise level on the same weekday within the same month (3 control days), with adjustment for regional background air pollution and temperature. In addition, we will analyze various other time-windows of wind turbine noise: on the previous day and up to 4 days (lag 4) before stroke/MI hospitalization and the accumulated exposure over 5 days (lag 0 – 4) as well as exposure at daytime (07:00–19:00 h), evening (19:00–22:00 h) and night (22:00–07:00 h).

Sub-study 2: Long-term exposure to wind turbine noise and risk for cardiovascular disease

We will use the unique Danish registers on wind turbines, residential addresses, health and socioeconomic status to conduct the first prospective study ever on effect of long-term exposure of noise from wind turbines on risk for cardiovascular disease.

Study population and design

The study population will consist of the 10,000-15,000 people (above 25 years of age) in Denmark who lived in dwellings within a noise immission area of one or more wind turbines in 1980-2012 (described in sub-study 1) as well as additionally 30,000 unexposed persons living just outside the noise immission areas of wind turbines, giving us a total study population of 40,000-45,000 persons.

We will identify all dwellings with exposure to wind turbine noise as described in sub-study 1. We will then identify unexposed persons using GIS and the official Danish address database (OIS) to identify dwellings within an area of approximately three times the distance from the wind turbine as for the exposed dwellings (Figure 3).

We will then identify all persons (above 25 years of age) who have lived in dwellings within the outer geographical circle (Figure 3) in the period between 1980 and 2012 by linking the addresses with the Danish civil registration system, thereby retrieving their unique personal identification number (33).

Similarly to the exposed population, we will only include persons who have lived at least one year at these addresses. By choosing this strategy for selection of an unexposed population, we will identify a population that are more similar to the exposed population than if we had chosen a random sample of 30,000 Danes.

We will determine the mobility of both the exposed and the unexposed population and based on this set up a strategy for handling moving out of the geographical areas, e.g. if the mobility is low we will exclude all cohort members who lived less than 80 % of the follow-up time in residences within the included geographical areas.

Information on hospitalization for cardiovascular disease (ICD10: I00-I99) in the period from 1977 to 2012 for these 40,000-45,000 persons will subsequently be found by linking their personal identification number to the Danish National Patient Registry (34). Furthermore, information on death will be collected from the Danish register of causes of death (35). We will exclude all participants hospitalized with cardiovascular disease before moving into dwellings within the geographical circles used in this study. We estimate that approximately 9,000 persons will be hospitalized with incident cardiovascular disease during the study period based on information on number of yearly incident cardiovascular events (www.hjerteforeningen.dk). We will generate sub-groups of cardiovascular diseases to be used in the statistical analyses: MI (ICD10: I21.0-I21.9; approximately 1,500 cases), atrial fibrillation and flutter (ICD10: I48; approximately 3,000 cases), heart failure (ICD10: I50; approximately 1,000 cases) and stroke (ICD10: I61, I63 and I64; approximately 2,000 cases).

We will use the wind turbine noise exposure data modeled for sub-study 1, and calculate 1- and 5-years averages for each person (see statistical section). We will investigate effects of whole day exposure as well as for nighttime exposure.

To be able to adjust all statistical analyses for socioeconomic status we will link the personal identification number of all population members to Statistic Denmark (36) to collect information on level of education, individual and household income, affiliation to the work market and cohabiting status.

Exposure to road traffic noise and/or air pollution could potentially confound the results. We have previously developed a GIS road network with traffic data for the period 1960–2005 (37). We will use this database together with the geocode of all dwellings in the study to derive three variables indicating the amount of traffic near each dwelling: distance to a road with a traffic density > 10,000 vehicle/day (major road), and two variables summarizing the total amount of kilometers driven by vehicles within 200 m and 500 m, respectively, of the residence each day as the product of street length and traffic density added up for all street lines within a 200/500 m circle around the address.




Figure 3 Identification of the unexposed population.

Statistical analyses

Statistical analyses will be based on a Cox proportional hazards model with age as the underlying time, which ensures comparison of individuals of the same age (38). We will use left truncation at age of enrolment into the population (January 1st 1985 (to allow for calculation of 5 year exposure preceding event), moving into an area close to a wind turbine (figure 3) or setting up of new wind turbine), so that people will be considered at risk from enrolment into the cohort, and end of follow-up at the age at diagnosis of a cardiovascular disease (event), death or December 31th 2012, whichever came first.

Exposure to wind turbine noise will be modeled as time-weighted averages the preceding 1- and 5-years at a given age. These exposures (1- and 5-years) will be entered as time-dependent variables into the statistical risk model, thus for each incident cardiovascular event recalculating exposure for all population members at exactly the same age as the case and at risk at the time of the hospitalization for cardiovascular disease. Time-weighted averages will be calculated based on 1) whole day exposure and 2) only nighttime exposure.

Estimates will be calculated crude and adjusted for a priori defined potential confounders: sex, level of education, calendar year, individual and household income, affiliation to the work market and cohabiting status. We will also adjust for distance to major road and traffic load within 200 m and 500 m of the residence as proxies for air pollution and road traffic noise.

We have calculated the power of this study based on a population of 40,000 where one third of the population is exposed. We have a statistical power of >90 % to detect associations of 1.10, 1.15 and 1.20 for all cardiovascular diseases (9,000 cases), atrial fibrillation and flutter (3,000 cases) and MI (1,500 cases), respectively. Proc power, SAS version 9.2, was used for these calculations.

SIGNIFICANCE AND DISSEMINATION

Exposure to wind turbine noise is suspected of affecting health and concern for this is increasing among people living close to wind turbines. However, very limited scientific data exists on effect of noise from wind turbines and risk for cardiovascular disease or other major diseases. This study will therefore be among the first contributing to answer the question of whether noise from wind turbines affects the risk of a major disease and has the potential to contribute in providing guidance on regulations for human habitation close to wind turbines.

Results will be published in international peer-reviewed journals and presented to the general population through national media and popular science articles.

FEASIBILITY

The Danish Cancer Society research group members have many years of experience in conducting environmental epidemiology research, and have during the last years specialized in investigating health effects of noise. We expect no problems in collecting and generating data and results for this study. Information on all Danish wind turbines is readily available from The Danish Energy Agency and Energinet.dk. We have extensive experience in collecting, managing and analyzing data from the national registers to be used in this study and expect no difficulties in gaining access to these registers or in obtaining permission from the Danish Data Protection Agency. We also have extensive experience in wind

speed and traffic modelling and the applicant has already been in contact with two acoustical consultant agencies interested in modelling the wind turbine noise for the project. Many of the research group members have successfully collaborated in several projects.

TIMELINES

The project runs for 24 months:

- Month 1: The Danish Cancer Society and DTU Wind Energy will meet with at least two acoustical consultant agencies to discuss and plan the wind turbine noise modeling. Based on this one consultant will be selected and a contract written and signed.
The Danish Cancer Society will apply the Danish Data Protection Agency and national registers (OIS, Statistic Denmark and 'Statens Serum Institut') for permissions for use of data for the study.
- Month 2-3: The consultant agency will generate a model for noise immission from all types of wind turbines to be used for identifying the exposed Danish dwellings.
- Month 2-6: DTU Wind Energy will model wind speed and direction time series at each wind turbine location.
- Month 4-5: Based on the modeled noise immission data the Danish Cancer Society will use the Danish address database (OIS) to identify addresses of all Danish dwellings exposed to wind turbine noise as well as unexposed dwellings for sub-study 2. Subsequently, these addresses will be linked to the Danish civil registration system, followed by linkage to Danish National Patient Registry, Danish register of causes of death and Statistic Denmark.
- Month 7-9: Based on wind modeling data and addresses of all exposed Danish dwellings the consultant agency will model wind turbine noise for all exposed dwellings.
Department of Environmental Science, Aarhus University, will generate traffic proxies and obtain information on historical background air pollution and temperature measurements.
- Month 10-15: The Danish Cancer Society will conduct the statistical analyses of sub-study 1.
- Month 16-20: The Danish Cancer Society will conduct the statistical analyses of sub-study 2.
- Month 16-24: The Danish Cancer Society will write scientific papers with input from all other research group members.

THE RESEARCH GROUP

The highly interdisciplinary research team will be headed by the applicant Mette Sørensen. The expertise of the team includes noise exposure, wind energy, register-based health research, use of GIS and biostatistics:

Mette Sørensen, PhD, senior researcher in the environment and cancer research group at the Danish Cancer Society, has many years of experience in environmental epidemiology. Since 2008 her research has focused on health effects of traffic noise, which among others have resulted in publications that as the first ever showed exposure to road traffic noise to be associated with risk for stroke and diabetes (3, 39). In March 2012 she received a 5-year starting grant from the European Research Council to investigate health consequences of noise exposure from road traffic (QUIET).

Ole Raaschou-Nielsen, PhD, head of the environment and cancer research group at the Danish Cancer Society, has many years of experience in health effects of traffic pollution, and participates in designing the study, choosing consultant agency, dialog with consultant agency etc.

Rikke Nordsborg, PhD student in the environment and cancer research group at the Danish Cancer Society. Rikke mastered in geography and has extensive experience in the use of GIS. She will be responsible for identification of dwellings in the vicinity of wind turbines using GIS, and for identifying persons living in the dwellings during the study period and link them to national registry data.

Alfredo Peña, PhD, senior scientist at DTU Wind Energy has extensive experience in wind resource assessment, wind power meteorology and meso-scale modeling. He will be responsible for providing the simulated wind speed and direction time series at the wind turbine locations.

All statistical analyses will be conducted by a **statistician/epidemiologist** who will be employed at the Danish Cancer Society.

Responsible for the modeling of wind turbine noise will be a **consultant agency** with extensive expertise within the area of wind turbine noise. To ensure the best value for money at least two consultant agencies will be approached.

Matthias Ketzel, PhD Senior Scientist, Department of Environmental Science, Aarhus University has extensive experience in air pollution modeling and will be responsible for generating traffic proxies and historical background concentrations of air pollution.

REFERENCES

1. Babisch W. Transportation noise and cardiovascular risk: updated review and synthesis of epidemiological studies indicate that the evidence has increased, 2006. *Noise Health* 8:1-29
2. Babisch W. Road traffic noise and cardiovascular risk, 2008. *Noise.Health* 10:27-33
3. Sorensen M, Hvidberg M, Andersen ZJ, et al. Road traffic noise and stroke: a prospective cohort study, 2011. *Eur Heart J* 32:737-744
4. Sorensen M, Andersen ZJ, Nordsborg RB, et al. Road traffic noise and incident myocardial infarction: a prospective cohort study, 2012. *PlosONE* 7:e39283
5. World Health Organization Regional Office for Europe. 2009. Night noise guidelines for Europe
6. Lusk SL, Gillespie B, Hagerty BM, Ziemba RA. Acute effects of noise on blood pressure and heart rate, 2004. *Arch.Environ.Health* 59:392-399
7. Wagner J, Cik M, Marth E, et al. Feasibility of testing three salivary stress biomarkers in relation to naturalistic traffic noise exposure, 2010. *Int.J Hyg.Environ Health* 213:153-155
8. Steptoe A, Hamer M, Chida Y. The effects of acute psychological stress on circulating inflammatory factors in humans: a review and meta-analysis, 2007. *Brain Behav.Immun.* 21:901-912
9. Haralabidis AS, Dimakopoulou K, Velonaki V, et al. Can exposure to noise affect the 24 h blood pressure profile? Results from the HYENA study, 2010. *J Epidemiol.Community Health*
10. Griefahn B, Brode P, Marks A, Basner M. Autonomic arousals related to traffic noise during sleep, 2008. *Sleep* 31:569-577
11. Miedema HM, Vos H. Associations between self-reported sleep disturbance and environmental noise based on reanalyses of pooled data from 24 studies, 2007. *Behav.Sleep Med.* 5:1-20
12. Ekstedt M, Akerstedt T, Soderstrom M. Microarousals during sleep are associated with increased levels of lipids, cortisol, and blood pressure, 2004. *Psychosom.Med.* 66:925-931
13. Stamatakis KA, Punjabi NM. Effects of sleep fragmentation on glucose metabolism in normal subjects, 2010. *Chest* 137:95-101
14. Wright CE, Erlich J, Valdimarsdottir HB, Bovbjerg DH. Poor sleep the night before an experimental stressor predicts reduced NK cell mobilization and slowed recovery in healthy women, 2007. *Brain Behav.Immun.* 21:358-363
15. Hui L, Hua F, Diandong H, Hong Y. Effects of sleep and sleep deprivation on immunoglobulins and complement in humans, 2007. *Brain Behav.Immun.* 21:308-310
16. Janssen SA, Vos H, Eisses AR, Pedersen E. A comparison between exposure-response relationships for wind turbine annoyance and annoyance due to other noise sources, 2011. *J.Acoust.Soc.Am.* 130:3746-3753
17. Pedersen E, Waye KP. Perception and annoyance due to wind turbine noise--a dose-response relationship, 2004. *J.Acoust.Soc.Am.* 116:3460-3470

18. Pedersen E, Persson WK. Wind turbine noise, annoyance and self-reported health and well-being in different living environments, 2007. *Occup. Environ. Med.* 64:480-486
19. Pedersen E. Health aspects associated with wind turbine noise - Results from three field studies, 2011. *Noise Control Engineering Journal* 59:47-53
20. Shepherd D, McBride D, Welch D, Dirks KN, Hill EM. Evaluating the impact of wind turbine noise on health-related quality of life, 2011. *Noise. Health* 13:333-339
21. Bakker RH, Pedersen E, van den Berg GP, Stewart RE, Lok W, Bouma J. Impact of wind turbine sound on annoyance, self-reported sleep disturbance and psychological distress, 2012. *Sci. Total Environ.* 425:42-51
22. Nissenbaum MA, Aramini JJ, Hanning CD. Effects of industrial wind turbine noise on sleep and health, 2012. *Noise. Health* 14:237-243
23. McColl BW, Allan SM, Rothwell NJ. Systemic infection, inflammation and acute ischemic stroke, 2009. *Neuroscience* 158:1049-1061
24. Biasucci LM, Leo M, De Maria GL. Local and systemic mechanisms of plaque rupture, 2008. *Angiology* 59:73S-76S
25. Søndergaard B, Henningsen P. Generelle data om støjen fra ældre vindmøller, 2011. Miljøstyrelsen Miljøprojekt Nr. 1398:
26. Søndergaard B, Plovsing B, Sørensen T. Validation of the Nord2000 propagation model for use on wind turbine noise, 2009. PSO-07 F&U project no.7389 Final report:1-53
27. Miljøministeriet. Bekendtgørelse om støj fra vindmøller, 2012. <https://www.retsinformation.dk/Forms/R0710.aspx?id=139658>
28. Hahmann AN, Lange J, Pena A, Hasager CB. The NORSEWInD numerical wind atlas for the South Baltic, 2012. DTU Wind Energy. DTU Wind Energy E, no 0011(EN).
29. Pena A, Gryning S-E, Hahmann AN. Observations of the atmospheric boundary layer height under marine upstream flow conditions at a coastal site, 2013. *J. Geophysical Research* 118:1924-1940
30. Marinelli M, Maule P, Hahmann AN, Isleifsson FR, Nørgård P, Cutululis N.A. Wind and photovoltaic large scale regional models for hourly production evaluation, IEEE transaction on sustainable energy, 2013. In preparation
31. Skamarock WC, Klemp JB, Dudhia J, et al. A description of the advanced research WRF version 3, 2008. NCAR/TN-475+STR Mesoscale and microscale meteorology division, National Center for Atmospheric Research, Boulder, Colorado, USA, 113pp:
32. Hahmann AN, Rostkier-Edelstein D, Warner TT, et al. A reanalysis system for the generation of mesoscale climatographies, 2010. *J Applied Meteorol Climatol* 49:954-972
33. Pedersen CB. The Danish Civil Registration System, 2011. *Scand. J. Public Health* 39:22-25
34. Lyng E, Sandegaard JL, Rebolj M. The Danish National Patient Register, 2011. *Scand. J. Public Health* 39:30-33
35. Helweg-Larsen K. The Danish Register of Causes of Death, 2011. *Scand. J. Public Health* 39:26-29
36. Thygesen LC, Daasnes C, Thaulow I, Bronnum-Hansen H. Introduction to Danish (nationwide) registers on health and social issues: structure, access, legislation, and archiving, 2011. *Scand. J. Public Health* 39:12-16

37. Jensen SS, Hvidberg M, Pedersen J, et al. GIS-based national street and traffic data base 1960-2005 (In Danish with English summary), 2009. National Environmental Research Institute, Aarhus University, Roskilde NERI Technical Report No. 678:
38. Thiebaut AC, Benichou J. Choice of time-scale in Cox's model analysis of epidemiologic cohort data: a simulation study, 2004. *Stat.Med.* 23:3803-3820
39. Sorensen M, Andersen ZJ, Nordsborg RB, et al. Long-term exposure to road traffic noise and incident diabetes: a cohort study, 2013. *Environ.Health Perspect.* 121:217-222