



Pesticide Residues in Food and Feed

PESTICIDE DETAILS

223 Thiacloprid

Functional Class

☐ Insecticide

Maximum Residue Limits for Thiacloprid

Commodity	MRL	Year of Adoption	Symbols	Note
Almond hulls	10 mg/Kg	2007		
Berries and other small fruits	1 mg/Kg	2007		
Cotton seed	0.02 mg/Kg	2007	(*)	
Cucumber	0.3 mg/Kg	2007		
Edible offal (mammalian)	0.5 mg/Kg	2007		
Egg plant	0.7 mg/Kg	2007		
Eggs	0.02 mg/Kg	2007	(*)	
Kiwifruit	0.2 mg/Kg	2007		
Meat (from mammals other than marine mammals)	0.1 mg/Kg	2007		
Melons, except watermelon	0.2 mg/Kg	2007		
Milks	0.05 mg/Kg	2007		
Mustard seed	0.5 mg/Kg	2007		
Peppers, Sweet (including pimento or pimiento)	1 mg/Kg	2007		
Pome fruits	0.7 mg/Kg	2007		
Potato	0.02 mg/Kg	2007	(*)	
Poultry meat	0.02 mg/Kg	2007	(*)	
Poultry, Edible offal of	0.02 mg/Kg	2007	(*)	
Rape seed	0.5 mg/Kg	2007		
Rice	0.02 mg/Kg	2007	(*)	
Squash, summer	0.3 mg/Kg	2007		
Stone fruits	0.5 mg/Kg	2007		
Tomato	0.5 mg/Kg	2007		
Tree nuts	0.02 mg/Kg	2007		
Watermelon	0.2 mg/Kg	2007		
Wheat	0.1 mg/Kg	2007		
Wheat straw and fodder, Dry	5 mg/Kg	2007		
Winter squash	0.2 mg/Kg	2007		

(*) At or about the limit of determination.

Other information

JMPR-related information

ADI/PTDI 0-0.01 mg/kg body weight -2006

Residue definition For compliance with MRLs and estimation of dietary intake in plant and animals commodities: thiacloprid. The residue is not fat soluble.

printer-friendly version

CODEX Pesticides Home [Commodities](#) [Pesticides](#) [Search](#) [Functional Classes](#)
Glossary

© FAO and WHO 2013
