



Pesticide Residues in Food and Feed

PESTICIDE DETAILS

213 Trifloxystrobin

Functional Class

☐ Fungicide

Maximum Residue Limits for Trifloxystrobin

Commodity	MRL	Year of Adoption	Symbols	Note
Almond hulls	3 mg/Kg	2006		
Asparagus	0.05 mg/Kg	2013	(*)	
Banana	0.05 mg/Kg	2006		
Barley	0.5 mg/Kg	2006		
Barley straw and fodder, Dry	7 mg/Kg	2006		
Brussels sprouts	0.1 mg/Kg	2006		
Cabbages, Head	0.5 mg/Kg	2006		
Carrot	0.1 mg/Kg	2006		
Celery	1 mg/Kg	2006		
Citrus fruits	0.5 mg/Kg	2006		
Citrus pulp, Dry	1 mg/Kg	2006		
Dried grapes (=currants, raisins and sultanas)	5 mg/Kg	2006		
Egg plant	0.7 mg/Kg	2013		
Eggs	0.04 mg/Kg	2006	(*)	
Flowerhead brassicas (includes Broccoli, Chinese and Cauliflower)	0.5 mg/Kg	2006		
Fruiting vegetables, Cucurbits	0.3 mg/Kg	2006		
Grapes	3 mg/Kg	2006		
Hops, Dry	40 mg/Kg	2006		
Kidney of cattle, goats, pigs and sheep	0.04 mg/Kg	2006	(*)	
Leek	0.7 mg/Kg	2006		
Lettuce, Head	15 mg/Kg	2013		
Liver of cattle, goats, pigs & sheep	0.05 mg/Kg	2006		
Maize	0.02 mg/Kg	2006		
Maize fodder (dry)	10 mg/Kg	2006		
Meat (from mammals other than marine mammals)	0.05 mg/Kg	2006	(fat)	
Milks	0.02 mg/Kg	2006	(*)	
Olive oil, Refined	1.2 mg/Kg	2013		
Olive oil, Virgin	0.9 mg/Kg	2013		
Olives	0.3 mg/Kg	2013		
Papaya	0.6 mg/Kg	2013		
Peanut	0.02 mg/Kg	2006	(*)	
Peanut fodder	5 mg/Kg	2006		
Peppers, Sweet (including pimento or pimiento)	0.3 mg/Kg	2006		
Pome fruits	0.7 mg/Kg	2006		
Potato	0.02 mg/Kg	2006	(*)	
Poultry meat	0.04 mg/Kg	2006	(*) (fat)	
Poultry, Edible offal of	0.04 mg/Kg	2006	(*)	
Radish	0.08 mg/Kg	2013		

Commodity	MRL	Year of Adoption	Symbols	Note
 Radish leaves (including radish tops)	15 mg/Kg	2013		
 Rice	5 mg/Kg	2006		
 Rice bran, Unprocessed	7 mg/Kg	2006		
 Rice straw and fodder, Dry	10 mg/Kg	2006		
 Stone fruits	3 mg/Kg	2006		
 Strawberry	1 mg/Kg	2013		
 Sugar beet	0.05 mg/Kg	2006		
 Sugar beet molasses	0.1 mg/Kg	2006		
 Sugar beet pulp, Dry	0.2 mg/Kg	2006		
 Tomato	0.7 mg/Kg	2006		
 Tree nuts	0.02 mg/Kg	2006	(*)	
 Wheat	0.2 mg/Kg	2006		
 Wheat bran, Unprocessed	0.5 mg/Kg	2006		
 Wheat straw and fodder, Dry	5 mg/Kg	2006		

(*) At or about the limit of determination.

(fat) (for meat) The MRL/EMRL applies to the fat of meat.

Other information

JMPR-related information

ADI/PTDI 0-0.04 mg/kg body weight -2004

Residue definition Definition of the residue for compliance with MRLs for plant commodities: trifloxystrobin; Definition of the residue for estimation of dietary intake of plant commodities: sum of trifloxystrobin and [(E,E)-methoxyimino-{2-[1-(3-trifluoromethylphenyl)ethylideneamino]oxymethyl}phenyl}acetic acid] (CGA 321113), expressed as trifloxystrobin. Definition of the residue for compliance with MRLs and estimation of dietary intake in animal commodities: sum of trifloxystrobin and [(E,E)-methoxyimino-{2-[1-(3-trifluoromethylphenyl)ethylideneamino-oxymethyl}phenyl}acetic acid] (CGA 321113), expressed as trifloxystrobinThe residue is fat-soluble.

 **printer-friendly version**

[CODEX Pesticides Home](#) [Commodities](#) [Pesticides](#) [Search](#) [Functional Classes](#)
[Glossary](#)

© FAO and WHO 2013